ریدازیکس®

سدیم کلراید ۰/۶۵ درصد

قبل از مصرف فرآورده، بروشور را به دقت مطالعه کنید.

موارد مصرف فرآورده:

شکل فرآورده: قطره بینی

قطره ریدازیکس حاوی سدیم کلراید ۰/۶۵ درصد است و برای کمک به رفع گرفتگی بینی و پاکسازی مجاری بینی مورد استفاده قرار می گیرد. محلول سالین به شل شدن و خارج شدن ترشحات و مخاط اضافی، آلرژن‌ها، گرد و غبار و سایر عوامل تحریک‌کننده از مجاری بینی کمک می‌کند و موجب راحت‌تر شدن تنفس می گردد. همچنین با مرطوب نگه داشتن مخاط بینی، به کاهش خشکی بینی ناشی از هوای خشک یا سرد، شرایط محیطی و یا مصرف برخی داروها کمک می‌کند.

راهنمایی های عمومی:

ریدازیکس برای بیماری فعلی شما تجویز شده است. لذا از مصرف آن در موارد مشابه یا توصیه به دیگران جدا خودداری نمایید.

هشدارها:

•مصرف ریدازیکس در کودکان تنها با تشخیص پزشک مجاز است.

•در صورتی که دریوش ظرف قبلا باز شده است، از مصرف آن جدا خودداری نمایید.

موارد احتیاط و منع مصرف:

•در صورت وجود هرگونه حساسیت نسبت به سدیم کلراید یا هر یک از مواد جانبی ذکر شده در بخش مشخصات دارو با پزشک خود مشورت نمایید.

مصرف در بارداری و شیردهی:

•در صورت بارداری و یا شیردهی پیش از مصرف ریدازیکس از پزشک یا داروساز خود مشورت بخواهید. این فرآورده در دوره بارداری و شیردهی قابل استفاده می باشد.

تداخلات دارویی:

در صورت استفاده از سایر داروها، حتما پزشک خود را مطلع نمایید.

مقدار مصرف فرآورده:

مقدار مصرف را پزشک معین می‌کند، اما مقدار مصرف معمول برای بزرگسالان و کودکان بالای ۲ سال، روزانه ۳ تا ۴ مرتبه و هر بار ۲ تا ۳ قطره در هر سوراخ بینی که لازم باشد بچکانید.

نحوه صحیح مصرف فرآورده:

•پیش از مصرف قطره دستان خود را بشویید.

•قبل از استفاده، بینی خود را به آرامی با بازدم هوا تخلیه کنید.

•پس از باز کردن دریوش بطری، بطری را بین انگشت شست و انگشت وسط نگه داشته، سر را مطابق شکل زیر متمایل کرده و نوک قطره چکان را با فاصله در سوراخ بینی قرار دهید. به ارامی به انتهای بطری فشار وارد نموده و قطره را داخل سوراخ بینی بچکانید. سر را برای لحظاتی در همان وضعیت نگه دارید.

•دریوش بطری را بلافاصله بعد استفاده محکم ببندید.

در موارد استفاده برای کودکان قبل از مصرف مجاری بینی کودک را با استفاده از پوآر بینی از تجمع مخاط پاک کنید.



عوارض جانبی احتمالی:

هر فرآورده به موازات آثار درمانی ممکن است باعث بروز برخی عوارض ناخواسته گردد، اگرچه همه این عوارض در یک فرد دیده نمی‌شود.

بروز عوارض جانبی به دنبال استفاده از این فراورده نادر است، اما در صورت بروز هرگونه عارضه مصرف فرآورده را قطع و با پزشک، داروساز مشورت کنید.

شرایط نگهداری فرآورده:

•فرآورده را دور از نور و در دمای کمتر از ۳۰ درجه سانتی‌گراد نگهداری و از یخ‌زدگی محافظت نمایید.

•فرآورده را دور از دید و دسترس کودکان نگهداری نمایید.

مشخصات فرآورده:

هر ۱ میلی‌لیتر از محلول ریدازیکس قطره بینی حاوی ۶/۵ میلی‌گرم ماده موثره سدیم کلراید می‌باشد. سایر مواد جانبی شامل بنزالکونیوم کلراید، سدیم‌دی ادتات، سدیم‌دی هیدروژن فسفات دی هیدرات، دی‌سدیم هیدروژن فسفات انهیدروز، گلیسیرین، و آب قابل تزریق (water for Injection) می‌باشد. هر جعبه حاوی ۲۰ میلی‌لیتر محلول در یک بطری ۲۰ میلی‌لیتری به همراه قطره چکان و دریوش می‌باشد.

ساخت شرکت داروسازی نوآوری زیستی
البرز - ایران

INOCLON

RIDAZIX®

Sodium Chloride 0.65%

Read all of this leaflet carefully before you start using this product because it contains important information for you.
•Keep this leaflet. You may need to read it again.
•Always use this product exactly as described in this leaflet or as instructed by your doctor or pharmacist.
•If you need more information or advice, ask your pharmacist.
•This product has been prescribed for you only. Do not pass it on to others. It may harm them, even if their symptoms are the same as yours.
•If you experience any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet.

What is in this leaflet:

1.What RIDAZIX nasal drop are and what they are used for
2.What you need to know before you use RIDAZIX nasal drop
3.How to use RIDAZIX nasal drop
4.Possible side effects
5.How to store RIDAZIX nasal drop
6.Contents of the pack and product description

1.What RIDAZIX nasal drop are and what they are used for

RIDAZIX nasal drops contain Sodium Chloride 0.65% and are intended to help relieve nasal congestion and cleanse the nasal passages.
The isotonic saline solution helps loosen and remove excess mucus, allergens, dust, and other irritants from the nasal passages, making breathing more comfortable.
RIDAZIX nasal drop also helps moisturize the nasal mucosa and relieve nasal dryness caused by dry or cold air, environmental conditions, or the use of certain medications.

2.What you need to know before you use RIDAZIX nasal drop

Caution

Do not use RIDAZIX nasal drop if:
•You are allergic (hypersensitive) to Sodium Chloride or any of the other ingredients in this product.
•The container is damaged or the solution appears abnormal.

Take special care with RIDAZIX nasal drop:
•RIDAZIX is generally suitable for children, adults, pregnant women, and elderly people when used as directed.
•For infants, especially newborns, or for prolonged use, consult your doctor or pharmacist before use.
•If nasal irritation, discomfort, or any unusual reaction occurs, discontinue use and consult your doctor or pharmacist.

Other product and RIDAZIX nasal drop:
•Tell your doctor or pharmacist if you are taking, have recently taken, or might take any other medicines.
Pregnancy, breast-feeding, and fertility:
•This product is safe to use if you are pregnant or breastfeeding.

3.How to use RIDAZIX nasal drop

Always use this product exactly as described in this leaflet or as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

How much to use
Adults and children over 2 years of age: Instill 2-3 drops into each nostril, 3-4 times daily, or directed by a physician.

Instructions for use in the nose
1.Wash your hands well before use.
2.Blow your nose gently.
3.Drop the required number of drops into each nostril.
4.The aim is to get the liquid to spread over all the inside surface of the nose, including the upper surface.
5.A good position is to lie on a bed with your head hanging back over the edge. Stay like this for two minutes after putting in the drops before getting up. This is to ensure that the liquid does not immediately run out of your nose or down the back of your throat but stays in the nasal cavity for a while.
6.Kneeling or bending forwards is an alternative, but it is harder to stay like this for two minutes after putting in the drops.
7.Replace the cap on the bottle after use.
8.Wash your hands after using the drops.

If you use more RIDAZIX nasal drop than you should:
If you are at all concerned that you have used too much of this product, contact your doctor or pharmacist.

If you forget to use RIDAZIX nasal drop:
Use required drops as soon as you remember and then go back to your regular dosing routine. However, if it is almost time for your next dose, skip the missed dose and go back to your regular dosing routine.

4.Possible side effects

Like all medications, this drug may cause some unwanted side effects, although not everyone experiences them. The occurrence of adverse effects with this product is rare; however, discontinue use and consult a doctor, pharmacist, or medical center if you experience any.

5.How to store RIDAZIX nasal drop:

•RIDAZIX nasal drop in the original container away from direct light.
•Store below 30 °C.
•Keep this product in a safe place where children cannot see or reach it.
•Throw away the bottle after 4 weeks of first opening.

6.Contents of the pack and product description

What RIDAZIX nasal drop:
The active substance is Sodium chloride 0.65% w/v.
The other ingredients are benzalkonium chloride, disodium edetate, Glycerin, Disodium hydrogen phosphate anhydrous, Sodium dihydrogen phosphate dihydrate and water for injection.

What RIDAZIX nasal drop looks like and the contents of the pack:
RIDAZIX nasal drop is a clear and colorless liquid supplied in a 20 mL bottle with a screw cap.

Name:	Sodiumchloride nasal drops leaflet
Edit:	01
Date:	1405.07.25
Size:	140x123mm

290c

287c