ریدازیکس®

سدیم کلراید ۰/۶۵ درصد

قبل از مصرف فرآورده، بروشور را به دقت مطالعه کنید.

موارد مصرف فرآورده:

دسته دارویی: شست و شو دهنده و مرطوب کننده مخاط

شکل دارویی: اسپری بینی

جهت تسکین احتقان، گرفتگی و خشکی بینی ناشی از خشکی هوا، سرماخوردگی، آلرژی، سینوزیت، حساسیت های خفیف یا مصرف بیش از حد داروهای ضد احتقان.

مکانیسم عملکرد ریدازیکس، کاهش ویسکوزیته مخاط بینی است که در اثر خاصیت اسمزی عروق بینی ایجاد می‌شود.

راهنمایی‌های عمومی:

ریدازیکس برای بیماری فعلی شما تجویز شده است. لذا از مصرف آن در موارد مشابه یا توصیه به دیگران جدا خودداری نمایید.

هشدارها:

•این فرآورده فقط با نسخه پزشک قابل تهیه است.

•مصرف ریدازیکس در کودکان تنها با تشخیص پزشک مجاز است.

•در صورتی که دریوش ظرف قبلا باز شده است، از مصرف آن جدا خودداری نمایید.

موارد احتیاط و منع مصرف:

•در صورت وجود هرگونه حساسیت نسبت به سدیم کلراید یا هر یک از مواد جانبی ذکر شده در بخش مشخصات فرآورده با پزشک خود مشورت نمایید.

مصرف در بارداری و شیردهی:

•مصرف ریدازیکس در دوران بارداری یا شیردهی بی‌خطر است.

تداخلات دارویی:

در صورت استفاده از سایر داروها، حتما پزشک خود را مطلع نمایید.

مقدار مصرف فرآورده:

مقدار مصرف هر فرآورده را پزشک معین می‌کند، اما مقدار مصرف معمول ریدازیکس ۲-۱ پاف هر ۴ ساعت یک‌بار در هر سوراخ بینی است.


نحوه صحیح مصرف فرآورده:

•پیش از مصرف اسپری دستان خود را بشویید.

•قبل از استفاده، بینی خود را به آرامی با بازدم هوا تخلیه کنید.

•پس از باز کردن دریوش اسپری، برای اطمینان از عملکرد پمپ، آن را دور از چشم و در هوا اسپری کنید تا بتوانید ذرات ریز را ببینید.

•اسپری را بین انگشت اشاره و انگشت وسط نگه داشته، سر را کمی به جلو خم کنید و به آرامی نفس بکشید. نوک اسپری را در سوراخ بینی قرار دهید، با دست سوراخ دیگر بینی را مسدود نمایید و بدون تماس با مخاط بینی اسپری را فشار دهید. نفس عمیق بکشید و از وارد شدن دارو به سوراخ بینی اطمینان حاصل نمایید.



•در صورت تماس نوک اسپری با مخاط بینی، نوک اسپری را با دستمال تمیز پاک نمایید.

•دریوش اسپری را بلافاصله پس از استفاده محکم ببندید.

استفاده بیش از یک نفر از این ظرف ممکن است باعث انتشار عفونت شود.

در صورت فراموشی هر نوبت از فرآورده، به محض یادآوری آن را مصرف کنید. ولی اگر تقریبا زمان نوبت بعدی فرا رسیده بود، مطابق روال قبل فرآورده را مصرف نموده و مقدار آن را دو برابر نکنید.

عوارض جانبی احتمالی:

هر فرآورده به موازات آثار درمانی ممکن است باعث بروز برخی عوارض ناخواسته گردد، اگرچه همه این عوارض در یک فرد دیده نمی‌شود.

در صورت بروز عوارض زیر یا عوارضی غیر از آنچه در برگه راهنما قید گردیده است فرآورده را قطع و با پزشک، داروساز و یا مراکز درمانی تماس بگیرید.

عطسه، سوزش، تحریک یا افزایش ترشحات بینی.

شرایط نگهداری فرآورده:

•فرآورده را دور از نور و در دمای کمتر از ۳۰ درجه سانتی‌گراد نگهداری و از یخ زدگی محافظت نمایید.

•فرآورده را دور از دید و دسترس کودکان نگهداری نمایید.

•حداکثر زمان نگهداری فرآورده پس از باز شدن چهار هفته است.

مشخصات دارو:

هر ۱ میلی‌لیتر از محلول اسپری بینی حاوی ۶/۵ میلی‌گرم ماده موثره سدیم کلراید می‌باشد. سایر مواد جانبی شامل بنزالکونیوم کلراید، سدیم دی‌هیدروژن فسفات دی‌هیدرات، دی‌سدیم هیدروژن فسفات انهیدروز، دی‌سدیم ادتات و آب قابل تزریق (water for injection) می‌باشد. هر جعبه حاوی ۲۰ میلی‌لیتر محلول در یک بطری ۲۰ میلی‌لیتری به همراه اسپری و دریوش می‌باشد.

ساخت شرکت داروسازی نوآوری زیستی
البرز - ایران

INOCLON

RIDAZIX®

Sodium Chloride 0.65%

Read all of this leaflet carefully before you start using this medicine because it contains important information for you.
•Keep this leaflet. You may need to read it again.
•Always use this medicine exactly as described in this leaflet or as instructed by your doctor or pharmacist.
•If you need more information or advice, ask your pharmacist.
•This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their symptoms are the same as yours.
•If you experience any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet.

What is in this leaflet:

1.What RIDAZIX nasal sprays are and what they are used for
2.What you need to know before you use RIDAZIX nasal sprays
3.How to use RIDAZIX nasal sprays
4.Possible side effects
5.How to store RIDAZIX nasal sprays
6.Contents of the pack and product description

1.What RIDAZIX nasal sprays are and what they are used for

Category: Nasal irrigant

RIDAZIX nasal sprays contain sodium chloride, a salt solution that treats dry or irritated nasal passages. RIDAZIX nasal sprays is used for treating blocked, dry or irritated nasal passages caused by cold and flu, sinusitis, allergy; including hayfever and post-nasal surgery. It works by rinsing and moisturizing the nasal passages.

For infants and young children with stuffy nose who cannot empty their noses, use of the product makes the mucus easier to remove with a nasal bulb syringe.

2.What you need to know before you use RIDAZIX nasal sprays

Caution

Do not use RIDAZIX nasal sprays if you are:

•Allergic (hypersensitive) to sodium chloride or any of the other ingredients in this medicine.

Take special care with RIDAZIX nasal sprays:

•Usually safe for children, pregnant women and elderly, but it is best to confirm with your doctor or pharmacist for infants or long-term use.

Other medicines and RIDAZIX nasal sprays:

•Tell your doctor or pharmacist if you are taking, have recently taken, or might take any other medicines.

Pregnancy, breast-feeding, and fertility:

•This product is safe to use if you are pregnant or you are breast-feeding a baby.

3.How to use RIDAZIX nasal sprays

Always use this medicine exactly as described in this leaflet or as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

How much to use:

Children and adults: Apply 1-2 puffs in each nostril every 4 hours or as directed by a doctor. May use a tissue to remove excess mucus.

Instructions for use in the nose

1.Wash your hands well before use.
2.Gently blow your nose to clear it of mucus before using the medicine.
3.Remove the cap. Shake the bottle. The first time you use the pump spray each day, you may have to "prime" it. Do this by squirting it a few times into the air until a fine mist comes out.
4.Tilt your head forward slightly. Breathe out slowly.
5.Hold the pump bottle with your thumb at the bottom and your index and middle fingers on top. Insert the canister tip in your nose, aiming the tip toward the back of your head. Use a finger on your other hand to close your nostril on the side not receiving the medicine.
6.Squeeze the pump as you begin to breathe in slowly through your nose. Repeat these steps for the other nostril. If you're using more than one spray in each nostril, follow all these steps again.
7.Try not to sneeze or blow your nose just after using the spray.
8.When you have finished, wipe the nozzle so it is clean and then put the cap back on.

For nasal use only. Administer as often as required one bottle should be used only by the same person as the use of this dispenser by more than one person may spread infection.

If you use more RIDAZIX nasal sprays than you should:

If you are at all concerned that you have used too much of this medicine contact your doctor or pharmacist.

If you forget to use RIDAZIX nasal sprays:

Use a single puff as soon as you remember and then go back to your regular dosing routine. However, if it is almost time for your next dose, skip the missed dose and go back to your regular dosing routine. Do not use a double dose to make up for the ones missed.

4.Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody experiences them.
If you experience any of the following side effects you should tell your doctor immediately. These are symptoms of an allergic reaction. Get emergency medical help if you have signs of an allergic reaction: hives, difficult breathing, or swelling of your face, lips, tongue, or throat.
Side effects usually do not occur with use of this product. However, if the inside of your nose is very dry and irritated, stinging may occur. If this effect lasts or gets worse, tell your doctor or pharmacist promptly.

Common side effects of RIDAZIX nasal sprays:

•A very serious allergic reaction to this drug is rare.
•If any of the side effects get serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

5.How to store RIDAZIX nasal sprays

•Store RIDAZIX nasal sprays in the original container away from direct light. Store below 30 °C.
•Keep this medicine in a safe place where children cannot see or reach it.
•Throw away the bottle after 4 weeks of first opening.

6.Contents of the pack and product description

What RIDAZIX nasal sprays contain:

The active substance is sodium chloride 0.65% w/v.
The other ingredients are benzalkonium chloride, sodium dihydrogen phosphate dihydrate, disodium hydrogen phosphate anhydrous, disodium edetate, and water for injection.

What RIDAZIX nasal sprays look like and the contents of the pack:

RIDAZIX nasal sprays is a clear and colorless liquid supplied in a 20 mL bottle with a spray cap.