تیر-آی® پرو

۰/۴ درصد

قبل از مصرف فرآورده، بروشور را به دقت مطالعه کنید.

موارد مصرف فرآورده:

دسته فرآورده: مرطوب‌کننده چشمی
شکل فرآورده: محلول لوبریکانت استریل چشمی

هیالورونات سدیم به صورت طبیعی در مخاط چشم وجود دارد، بنابراین این قطره با خصوصیات مشابه به مایع مخاطی چشم، سازگاری زیادی با چشم دارد. این ماده توانایی بالایی در نگهداری آب دارد و به همین علت برای مرطوب کردن سطح چشم بسیار مناسب است. همچنین موجب افزایش پایداری پرده اشکی و احساس بهبود در سطح چشم تحریک شده می‌شود. تحریکات چشمی معمولاً بر اثر عواملی مانند باد، تابش خورشید، هوای خشک سرد/گرم، آب‌شور، دود، نور شدید، استفاده طولانی‌مدت از کامپیوتر، جراحی‌های‌چشمی، التهاب ملتحمه (کونژنکتیویت) یا استفاده از لنزهای تماسی ایجاد می‌شوند.

هشدارها:

•در صورتی که دریوش ظرف قبلا باز شده است، از مصرف آن جدا خودداری نمایید.

موارد احتیاط و منع مصرف:

•در صورت وجود هرگونه حساسیت نسبت به هیالورونات سدیم یا هر یک از مواد جانبی ذکر شده در بخش مشخصات فرآورده با پزشک خود مشورت نمایید.
•این فرآورده جهت مصرف چشمی می‌باشد. از مصرف فرآورده به صورت خوراکی خودداری شود.
•از تماس نوک قطره‌چکان با چشم یا سطوح دیگر خودداری نمایید.
•به دلیل احتمال تاری دید، بلافاصله پس از مصرف قطره تا برگشتن دید کامل، از رانندگی و کار با ماشین‌آلات بپرهیزید.
•از مصرف همزمان با لنزهای تماسی نرم خودداری نمایید.

مصرف در بارداری و شیردهی:

•در صورت بارداری و یا تصمیم برای باردار شدن پیش از مصرف تیر-آی‌پرو از پزشک یا داروساز خود مشورت بخواهید.

تداخلات فرآورده:

در صورت استفاده از سایر داروها، حتما پزشک خود را مطلع نمایید.
در صورت استفاده از پماد چشمی یا سایر قطره‌های چشمی، حداقل باید ۵ دقیقه فاصله زمانی بین مصرف دو فرآورده در نظر گرفته شود. پمادهای چشمی باید در مرحله آخر مصرف شوند.

مقدار مصرف فرآورده:

مقدار مصرف هر فرآورده را پزشک معین می‌کند، اما مقدار مصرف معمول تیر-آی‌پرو به شرح زیر است:
روزی ۳-۲ مرتبه هر بار یک قطره در چشم چکانده شود.

نحوه صحیح مصرف فرآورده:

•پیش از مصرف قطره دستان خود را بشویید.
•پس از باز کردن دریوش بطری، بطری را بین انگشت شست و انگشت وسط نگه‌داشته، کمی سر را به عقب متمایل کرده، با انگشت تمیز، پلک چشم را پایین کشیده و به آرامی به انتهای بطری فشار وارد نمایید. (مطابق شکل).



•قطره را در حفره به وجود آمده بین پلک و چشم بریزید. نوک بطری را نزدیک چشم برده اما مراقب باشید با سطح چشم، پلک و اطراف آن برخورد نکند زیرا احتمال آلوده شدن قطره وجود دارد.
•در حالیکه به سمت پایین نگاه می کنید چشم را به آرامی ببندید.
•به مدت ۳ تا ۵ دقیقه به کمک انگشت شست، گوشه چشم را به آرامی فشار دهید. چشم را بسته نگه‌دارید و در این فرصت داروی اضافی خارج شده از چشم را با دستمال پاک کنید.
•دریوش بطری را بلافاصله پس از استفاده محکم ببندید.

عوارض جانبی احتمالی:

هر فرآورده به موازات آثار درمانی ممکن است باعث بروز برخی عوارض ناخواسته گردد، اگرچه همه این عوارض در یک فرد دیده نمی‌شود. در صورت بروز عوارض زیر یا عوارضی غیر از آنچه در برگه راهنما قید گردیده است فرآورده را قطع و با پزشک داروساز و یا مراکز درمانی تماس بگیرید.
واکنش‌های آلرژیک از جمله سوزش ملایم، خارش، قرمزی و تاری دید ممکن است، بلافاصله بعد از مصرف رخ دهد که تا چند دقیقه پس از مصرف از بین می رود.

شرایط نگهداری:

•فرآورده را دور از نور و دمای کمتر از ۳۰ درجه سانتی‌گراد نگهداری و از یخزدگی محافظت نمایید.
•فرآورده را دور از دید و دسترس کودکان نگهداری نمایید.
•حداکثر زمان نگهداری فرآورده پس از باز شدن چهار هفته است.

مشخصات فرآورده:

هر ۱ میلی‌لیتر از محلول قطره استریل چشمی حاوی ۴ میلی‌گرم ماده موثره هیالورونات سدیم می‌باشد. سایر مواد جانبی شامل سدیم کلراید، پلی‌سوربات ۸۰، دی‌سدیم هیدروژن فسفات بدون آب، بنزالکونیوم کلراید، سدیم دی‌هیدروژن فسفات دی‌هیدراته، دی‌سدیم ادتات و آب قابل تزریق (water for injection) می‌باشد. هر جعبه حاوی ۱۰ میلی‌لیتر محلول در یک بطری ۱۰ میلی‌لیتری به همراه قطره چکان و دریوش استریل می‌باشد.

ساخت شرکت داروسازی نوآوری زیستی البرز - ایران

INOCLON

TEAR-I PRO®

Sodium hyaluronate 0.4%

Read all of this leaflet carefully before you start using this medicine because it contains important information for you.
•Keep this leaflet. You may need to read it again.
•Always use this medicine exactly as described in this leaflet or as instructed by your doctor or pharmacist.
•Ask your pharmacist if you need more information or advice.
•This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
•If you experience any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet.

What is in this leaflet:

1.What TEAR-I PRO® eye drops are and what they are used for.
2.What you need to know before you use Sodium TEAR-I PRO® drops.
3.How to use TEAR-I PRO® eye drops.
4.Possible side effects.
5.How to store TEAR-I PRO® eye drops.
6.Contents of the pack and product description.

1.What TEAR-I PRO® eye drops are and what they are used for

Category: Eye Lubricant

TEAR-I PRO® occurs naturally in the eye and is well tolerated. It binds water effectively,keeping the surface of the eye hydrated, lubricated, and protected. By stabilizing the tear film, it helps relieve dryness and irritation caused by factors such as wind, sun, dry air, smoke, computer use, surgery, conjunctivitis, or contact lens wear.

2.What you need to know before you use TEAR-I PRO® eye drops

Caution

Do not use TEAR-I PRO® eye drops if:

•You are allergic (hypersensitive) to sodium hyaluronate or any of the other ingredients in this medicine.
•If the container is damaged.
•Do not use with soft contact lenses simultaneously.

Other medicines and TEAR-I PRO® eye drops

•Tell your doctor or pharmacist if you are taking, have recently taken, or might take any other medicines.
•If you are using other eye drops or eye ointments, leave at least 5 minutes between each medicine. Eye ointments should be administered last.

Driving and using machines:

Your vision may be blurred for a short time after using TEAR-I PRO® eye drops. Make sure your sight is clear before driving or using machines.

Pregnancy, breastfeeding, and fertility

•If you are pregnant, or breastfeeding your baby, please consult your doctor, pharmacist or other healthcare professional for advice before using TEAR-I PRO®.

3.How to use TEAR-I PRO® eye drops

Always use this medicine exactly as described in this leaflet or as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

How much to use

Your doctor will determine the appropriate dosage, but the usual doses are as follows:

Use one drop in the affected eye(s) two to three times daily.

Instructions for use in the eye:

If you wear soft contact lenses, these must be removed during treatment of the eyes with TEAR-I PRO® drops.

1.Before opening a new bottle of TEAR-I PRO® drops, please check that the safety strip on the bottom of the out er cap is unbroken.
2.Twist the outer cap to break the seal
3.Wash your hands well before use.
4.Remove the outer cap. If you are putting in the drops yourself, you should use a mirror.
5.Tilt your head back and look up at the ceiling.
6.Pull the lower lid of the eye out to form a small pocket between your eyelid and your eye.
7.Hold the container between the thumb and middle finger of the outer hand, turn the bottle upside down near to your eye, try not to touch your eye with the nozzle.
8.Apply enough pressure to the container to release one to two drops, or as directed by your doctor.
9.Blink your eye a few times.
10.If you think that you have missed your eye, then insert another drop.
11.Repeat steps 5-10 in the other eye, if you have been told to use TEAR-I PRO® drops in both eyes.
12.Replace the outer cap on the bottle, trying not to touch the applicator tip with anything, including the eye or your fingers.
13.Wash your hands.

If you use more TEAR-I PRO® eye drops than you should:

It will not cause you any harm. If you are however worried, talk to your doctor or pharmacist. In the event of overdosage, consult your doctor or pharmacist. If neither is available, contact the nearest hospital.

TEAR-I PRO® eye drops are for external use only.

If you forget to use TEAR-I PRO® eye drops:

Use a single drop as soon as you remember and then go back to your regular dosing routine. However, if it is almost time for your next dose, skip the missed dose and go back to your regular dosing routine. Do not use a double dose to make up for the one missed.

If you stop using TEAR-I PRO® eye drops:

You should keep using TEAR-I PRO® drops for as long as your doctor has instructed. Your treatment is helping your condition, even if you have not noticed a difference in how you feel. Should you need to stop your treatment,

your doctor will advise you how to do this.

4.Possible side effects

Like all medications, this drug may cause some unwanted side effects, although not everyone experiences them. Discontinue use and consult a doctor, pharmacist, or medical center if you experience any of the following or other unexpected effects: blurred vision, eye irritation, burning sensation, eye pain, itching

5.How to store TEAR-I PRO® eye drops

•Store TEAR-I PRO® eye drops in the original container away from direct light.
•Store below 30°C.
•Keep this medicine in a safe place where children cannot see or reach it.
•Throw away the bottle after 4 weeks of first opening.

6.Contents of the pack and product description

What TEAR-I PRO® eye drops contain:
The active substance is sodium hyaluronate. Each mL contains 4mg of sodium hyaluronate. The other ingredients are, sodium chloride, polysorbate 80, disodium hydrogen phosphate anhydrous, benzalkonium chloride, sodium dihydrogen phosphate dihydrate, disodium edetate and water for injection.

What TEAR-I PRO® eye drops look like and the contents of the pack:

TEAR-I PRO® eye drops is a clear and colorless liquid supplied in a pack of 10 mL bottle with a screw cap.